

DEFINITIONS OF TNM

Definition of Primary Tumor (T)

T Category	Thickness	Ulceration status
TX: primary tumor thickness cannot be assessed (e.g., diagnosis by curettage)	Not applicable	Not applicable
T0: no evidence of primary tumor (e.g., unknown primary or completely regressed melanoma)	Not applicable	Not applicable
Tis (melanoma <i>in situ</i>)	Not applicable	Not applicable
T1	≤1.0 mm	Unknown or unspecified
T1a	<0.8 mm	Without ulceration
T1b	<0.8 mm 0.8–1.0 mm	With ulceration With or without ulceration
T2	>1.0–2.0 mm	Unknown or unspecified
T2a	>1.0–2.0 mm	Without ulceration
T2b	>1.0–2.0 mm	With ulceration
T3	>2.0–4.0 mm	Unknown or unspecified
T3a	>2.0–4.0 mm	Without ulceration
T3b	>2.0–4.0 mm	With ulceration
T4	>4.0 mm	Unknown or unspecified
T4a	>4.0 mm	Without ulceration
T4b	>4.0 mm	With ulceration

Definition of Regional Lymph Node (N)

Extent of regional lymph node and/or lymphatic metastasis		
N Category	Number of tumor-involved regional lymph node	Presence of in-transit, satellite, and/or microsatellite metastases
NX	Regional nodes not assessed (e.g., SLN biopsy not performed, regional nodes previously removed for another reason) Exception: pathological N category is not required for T1 melanomas, use cN.	No
N0	No regional metastases detected	No
N1	One tumor-involved node or in-transit, satellite, and/or microsatellite metastases with no tumor-involved nodes	
N1a	One clinically occult (i.e., detected by SLN biopsy)	No
N1b	One clinically detected	No
N1c	No regional lymph node disease	Yes
N2	Two or three tumor-involved nodes or in-transit, satellite, and/or microsatellite metastases with one tumor-involved node	

Extent of regional lymph node and/or lymphatic metastasis		
N Category	Number of tumor-involved regional lymph node	Presence of in-transit, satellite, and/or microsatellite metastases
N2a	Two or three clinically occult (i.e., detected by SLN biopsy)	No
N2b	Two or three, at least one of which was clinically detected	No
N2c	One clinically occult or clinically detected	Yes
N3	Four or more tumor-involved nodes or in-transit, satellite, and/or microsatellite metastases with two or more tumor-involved nodes, or any number of matted nodes without or with in-transit, satellite, and/or microsatellite metastases	
N3a	Four or more clinically occult (i.e., detected by SLN biopsy)	No
N3b	Four or more, at least one of which was clinically detected, or presence of any number of matted nodes	No
N3c	Two or more clinically occult or clinically detected and/or presence of any number of matted nodes	Yes

Definition of Distant Metastasis (M)

M Criteria		
M Category	Anatomic site	LDH level
M0	No evidence of distant metastasis	Not applicable
M1	Evidence of distant metastasis	See below
M1a	Distant metastasis to skin, soft tissue including muscle, and/or nonregional lymph node	Not recorded or unspecified
M1a(0)		Not elevated
M1a(1)		Elevated
M1b	Distant metastasis to lung with or without M1a sites of disease	Not recorded or unspecified
M1b(0)		Not elevated
M1b(1)		Elevated
M1c	Distant metastasis to non-CNS visceral sites with or without M1a or M1b sites of disease	Not recorded or unspecified
M1c(0)		Not elevated
M1c(1)		Elevated
M1d	Distant metastasis to CNS with or without M1a, M1b, or M1c sites of disease	Not recorded or unspecified
M1d(0)		Normal
M1d(1)		Elevated

Suffixes for M category: (0) LDH not elevated, (1) LDH elevated. No suffix is used if LDH is not recorded or is unspecified.

AJCC PROGNOSTIC STAGE GROUPS

Clinical (cTNM)

Clinical staging includes microstaging of the primary melanoma and clinical/radiologic/biopsy evaluation for metastases. By convention, clinical staging should be used after biopsy of the primary melanoma, with clinical assessment for regional and distant metastases. Note that pathological assessment of the primary melanoma is used for both clinical and pathological classification. Diagnostic biopsies to evaluate possible regional and/or distant metastasis also are included. Note there is only one stage group for clinical Stage III melanoma.

When T is...	And N is...	And M is...	Then the clinical stage group is...
Tis	N0	M0	0
T1a	N0	M0	IA
T1b	N0	M0	IB
T2a	N0	M0	IB
T2b	N0	M0	IIA
T3a	N0	M0	IIA
T3b	N0	M0	IIB
T4a	N0	M0	IIB
T4b	N0	M0	IIC
Any T, Tis	≥N1	M0	III
Any T	Any N	M1	IV

Pathological (pTNM)

Pathological staging includes microstaging of the primary melanoma, including any additional staging information from the wide-excision (surgical) specimen that constitutes primary tumor surgical treatment and pathological information about the regional lymph nodes after SLN biopsy or therapeutic lymph node dissection for clinically evident regional lymph node disease.

When T is...	And N is...	And M is...	Then the pathological stage group is...
Tis	N0	M0	0
T1a	N0	M0	IA
T1b	N0	M0	IA
T2a	N0	M0	IB
T2b	N0	M0	IIA
T3a	N0	M0	IIA
T3b	N0	M0	IIB
T4a	N0	M0	IIB
T4b	N0	M0	IIC
T0	N1b, N1c	M0	IIIB
T0	N2b, N2c, N3b or N3c	M0	IIIC
T1a/b–T2a	N1a or N2a	M0	IIIA

When T is...	And N is...	And M is...	Then the pathological stage group is...
T1a/b–T2a	N1b/c or N2b	M0	IIIB
T2b/T3a	N1a–N2b	M0	IIIB
T1a–T3a	N2c or N3a/b/c	M0	IIIC
T3b/T4a	Any N ≥N1	M0	IIIC
T4b	N1a–N2c	M0	IIIC
T4b	N3a/b/c	M0	IIID
Any T, Tis	Any N	M1	IV

Pathological Stage 0 (melanoma *in situ*) and T1 do not require pathological evaluation of lymph nodes to complete pathological staging; use cN information to assign their pathological stage.

r/yc/yp Classification

For additional information on recurrent/retreatment (r) and/or posttherapy/post neoadjuvant therapy (yc/yp) staging, please refer to Chapter 1.

REGISTRY DATA COLLECTION VARIABLES

1. Breslow tumor thickness (xx.x mm)
2. Primary tumor ulceration (yes/no)
3. Mitotic rate (whole number per square millimeter [mm²])
4. Microsatellites (pathologically detected satellites, not clinically apparent) (yes/no)
5. Tumor-infiltrating lymphocytes (absent, nonbrisk, or brisk)
6. Clark level of invasion (I–V)
7. Regression (yes/no)
8. Neurotropism (present or absent)
9. Lymphovascular invasion (present or absent)
10. In-transit and/or satellite metastasis (in-transit, satellite, both)
11. Regional lymph node clinically or radiologically detected (yes/no)
12. Microscopic confirmation of tumor metastasis in any regional lymph node that was clinically or radiologically detected (yes/no)
13. SLN biopsy performed (yes/no)
14. Number of nodes examined from sentinel node procedure (whole number)
15. Number of tumor-involved nodes from sentinel node procedure (whole number)
16. Sentinel node tumor burden (largest dimension of largest discrete deposit in xx.x mm)
17. ENE in any tumor-involved regional lymph node (sentinel or clinically detected) (present or absent)
18. Completion or therapeutic lymph node dissection performed (yes/no)